Smad3 (phospho-S204) polyclonal antibody

Catalog: AP0328 Host: Rabbit Reactivity: Human,Mouse,Rat

BackGround:

Smad proteins, the mammalian homologs of the Drosophila mothers against decapentaplegic (Mad), have been implicated as downstream effectors of GFβ/BMP signaling. Smad1 (also designated Madr1 or JV4-1) and Smad5 are effectors of BMP-2 and BMP-4 function, while Smad2 (also designated Madr2 or JV18-1) and Smad3 are involved in TGFβ and activin-mediated growth modulation. Smad4 (also designated DPC4) has been shown to mediate all of the above activities through interaction with various Smad family members. Smad6 and Smad7 regulate the response to activin/TGFβ signaling by interfering with TGFβ-mediated phosphorylation of other Smad proteins.

Product:

1 mg/ml in Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.2.

Molecular Weight:

~ 48, 55 kDa

Swiss-Prot:

P84022

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 100% (by SDS-PAGE).

Applications:

WB:1:500~1:1000

Storage&Stability:

Store at 4℃ short term. Aliquot and store at -20℃ long term. Avoid freeze-thaw cycles.

Specificity:

p-Smad3 (S204) polyclonal antibody detects endogenous levels of Smad3 protein only when phosphorylated at Ser204.

DATA:



Western blot (WB) analysis of p-Smad3 (S204) polyclonal antibody at 1:500 dilution

Lane1:HEK293T cell lysate treated with serum (20%, 30mins)

Lane2:sp2/0 cell lysate treated with serum (20%, 30mins)

Lane3:H9C2 cell lysate treated with serum (20%, 30mins)

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.
Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.
Email: info@bioworlde.com
Tel: 6123263284
Fax: 6122933841

Bioworld technology, co. Ltd.
Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.
Email: info@biogot.com
Tel: 0086-025-68037686
Fax: 0086-025-68035151