

PRODUCT DATA SHEET



Bioworld Technology, Inc.

Recombinant HB-EGF, Human

Catalog Number: BK0218-1mg

Source: CHO

Quantity: 1mg

Description:

Proheparin-binding EGF-like growth factor (HB-EGF), also known as DTR, DTS and HEGFL, is a member of the EGF family of mitogens. It is expressed in macrophages, monocytes, endothelial cells and muscle cells. HB-EGF signals through the EGF receptor to stimulate the proliferation of smooth muscle cells, epithelial cells and keratinocytes. Compared to EGF, HB-EGF binds the EGF receptor with higher affinity and is thus more mitogenic, probably due to its ability to bind to heparin and heparin sulfate proteoglycans. HB-EGF has been reported to act as a diphtheria toxin receptor, mediating endocytosis of the bound toxin.

Molecular Weight:

12-14 kDa, observed by reducing SDS-PAGE.

Purity:

> 95% as analyzed by SDS-PAGE and HPLC.

Biological Activity:

ED50 <0.5 ng/ml, measured in a cell proliferation assay using 3T3 cells.

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) pow-

der.

Formulation:

Lyophilized after extensive dialysis against PBS.

AA Sequence:

DLQEADLDLLRVTLSKPKQALATPNKEE-
HGKRKKKKGKGLGKKRDPCLRKYKDFCIH-
GECKYVKELRAPSCICHPGYHGERCHGLSL

Endotoxin:

< 0.2 EU/μg, determined by LAL method.

Reconstitution:

Reconstituted in ddH₂O or PBS at 100 μg/ml.

Storage:

Lyophilized recombinant Human HB-EGF remains stable up to 6 months at -80 °C from date of receipt. Upon reconstitution, Human HB-EGF should be stable up to 1 week at 4 °C or up to 2 months at -20 °C.

Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. For research use only.