

# FADD (N188) polyclonal antibody

Catalog: BS1399

Host: Rabbit

Reactivity: Human, Mouse, Rat

## **BackGround:**

In contrast to growth factors which promote cell proliferation, FAS ligand (FAS-L) and the tumor necrosis factors (TNFs) rapidly induce apoptosis. Cellular response to FAS-L and TNF is mediated by structurally related receptors containing a conserved "death domain" and belonging to the TNF receptor superfamily. TRADD, FADD and RIP are FAS/TNF-R1 interacting proteins that contain a death domain homologous region (DDH). TRADD (TNF-R1-associated domain) death and FADD (FAS-associated death domain) associate with the death domains of both FAS and TNF-R1 via their DDH regions. Overexpression of TRADD leads to NFkB activation and apoptosis in the absence of TNF. Overexpression of FADD causes apoptosis, which can be blocked by the cow pox protein CrmA, suggesting that FADD lies upstream of ICE and possibly other serine proteases. The receptor interacting protein, RIP, associates with FAS exclusively via its DDH and this association is abrogated in lpr mutants. Unlike TRADD and FADD, RIP contains a putative amino-terminal kinase domain.

#### **Product:**

1 mg/ml in Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

**Molecular Weight:** 

### ~ 28 kDa

**Swiss-Prot:** 

#### Q13158

**Purification&Purity:** 

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

**Applications:** 

WB: 1:500~1:1000

IHC: 1:50~1:200

**Storage&Stability:** 

#### Bioworld Technology, Inc.

 
 Add:
 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.

 Email:
 info@bioworlde.com


 Tel:
 6123263284

 Fax:
 6122933841
 Store at  $4 \,^{\circ}{\rm C}$  short term. Aliquot and store at  $-20 \,^{\circ}{\rm C}$  long term. Avoid freeze-thaw cycles.

#### **Specificity:**

FADD (N188) polyclonal antibody detects endogenous levels of FADD protein.

**DATA:** 



Western blot (WB) analysis of FADD (N188) polyclonal antibody at 1:500 dilution

Lane1:A549 whole cell lysate(40ug) Lane2:HCT116 whole cell lysate(40ug)

2

Lane3:Panc1 whole cell lysate(40ug)

Lane4:C6 whole cell lysate(40ug)

Lane5:MEF whole cell lysate(40ug)



Immunohistochemistry (IHC) analyzes of FADD (N188) pAb in paraf-

fin-embedded human lung carcinoma tissue at 1:100.

## Note:

For research use only, not for use in diagnostic procedure.

#### Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: info@biogot.com Tel: 0086-025-68037686 Fax: 0086-025-68035151



Bioworld Technology,Inc.

Bioworld Technology, Inc.		Bioworld technology, co. Ltd.	
Add:	1660 South Highway 100, Suite 500 St. Louis Park,	Add:	No 9, weidi road Qixia District Nanjing, 210046,
	MN 55416,USA.		P. R. China.
Email:	<u>info@bioworlde.com</u>	Email:	info@biogot.com
Tel:	6123263284	Tel:	0086-025-68037686
Fax:	6122933841	Fax:	0086-025-68035151