Bcl-X (S43) polyclonal antibody

Catalog:	BS1512	Host:	Rabbit	Reactivity:	Human,Mouse,Rat

BackGround:

Overexpression of Bax accelerates apoptotic death induced by cytokine deprivation in an IL-3-dependent cell line, and Bax also counters the death repressor activity of Bcl-2. Bcl-x, one of several additional proteins with sequence homology to Bcl-2, is expressed as Bcl-xL, a 233 amino acid protein with 43% sequence identity with Bcl-2 that suppresses cell death, and Bcl-xS, a shorter variant that is 178 amino acids in length and lacks a 63 amino acid region (amino acids 126-188) found in Bcl-xL and which functions as a dominant inhibitor of Bcl-2. A further apoptosis-inducing protein, Bad, dimerizes both with Bcl-xL and to a lesser extent with Bcl-2, thus displacing Bax and inducing apoptosis

Product:

1 mg/ml in Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

Molecular Weight:

~ 26 kDa

Swiss-Prot:

Q07817

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

**Applications:**

IHC: 1:50~1:200

IF: 1:50~1:200

Storage&Stability:

Store at 4 ℃ short term. Aliquot and store at -20 ℃ long term. Avoid freeze-thaw cycles.

Specificity:

BCL-X (S43) polyclonal antibody detects endogenous levels of BCL-XL protein, and also detects Isoform Bcl-XS (19 KDa).

DATA:

Immunohistochemistry (IHC) analyzes of Bcl-X (S43) pAb in paraffin-embedded human breast carcinoma tissue at 1:100.

Note:

For research use only, not for use in diagnostic procedure.