

c-Fms (E557) polyclonal antibody

Catalog: BS1627

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

c-Fms/CSF-1R is a transmembrane tyrosine kinase receptor belonging to the CSF1/PDGF receptor family. It is encoded by the c-Fms proto-oncogene and is expressed in mononuclear phagocytes, oocytes, decidual cells, trophoblastic cells and some myoblasts. It is important for growth and differentiation of myeloid cells and its function can be regulated by SLAP-2. c-Fms/CSF-1R is responsible for mediating all of the functions of M-CSF. M-CSF is a glycoprotein required for the proliferation and differentiation of mononuclear phagocytes, including osteoclasts. M-CSF has also been identified as an important mediator of the inflammatory response and can regulate the release of proinflammatory cytokines from macrophages.

Product:

1 mg/ml in Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

Molecular Weight:

Predicted band size : 108 kDa

Observed band size : 130 kDa

Swiss-Prot:

P07333

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

IHC: 1:50~1:200

IF: 1:50~1:200



Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

c-Fms (E557) polyclonal antibody detects endogenous levels of c-Fms protein.

DATA:



Immunohistochemistry (IHC) analyzes of c-Fms (E557) pAb in paraffin-embedded human brain carcinoma tissue at 1:50.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151