

Clock (E270) polyclonal antibody

Catalog: BS1873

Host: Rabbit

Reactivity: Human

BackGround:

Clock, a member of the basic-helix-loop-helix-PAS (bHLH-PAS) family of transcription factors, has also been identified as having circadian function. Mutations within the Clock gene have been shown to increase the length of the endogenous period and to cause a loss of rhythmicity of circadian oscillations. Clock contains a DNA-binding domain, a protein dimerization domain and a glutamine-rich C-terminal region, which indicates transactivation capabilities. It has been speculated that Clock may regulate circadian rhythmicity in combination with other proteins such as Per.

Product:

1 mg/ml in Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

Molecular Weight:

~ 100 kDa

Swiss-Prot:

O15516

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Clock (E270) polyclonal antibody detects endogenous levels of Clock protein.

DATA:




Western blot (WB) analysis of Clock (E270) pAb at 1:500 dilution

Lane1:A549 whole cell lysate(40ug)


Lane2:HEK293T whole cell lysate(40ug)

Lane3:A2780 whole cell lysate(40ug)



BS1873
Lot CJ36131

Immunohistochemistry (IHC) analyzes of Clock (E270) pAb in paraffin-embedded human colon carcinoma tissue at 1:50. showing Cytoplasm and Nucleus staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.



BS1873
Lot CJ36131

Immunohistochemistry (IHC) analyzes of Clock (E270) pAb in paraffin-embedded human colon carcinoma tissue at 1:50. showing Cytoplasm and Nucleus staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.

Note:

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151



PRODUCT DATA SHEET

Bioworld Technology, Inc.

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,
MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,
P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151