

EKLF (W318) polyclonal antibody

| Catalog: | BS2303 | Host: | Rabbit | Reactivity: Mouse,Rat |
|--|--|-------|-------------|--|
| | | | | |
| BackGround: | | | | Applications: |
| The Kruppel-type zinc finger transcription factors com- | | | | WB: 1:1000~1:5000 |
| prise a conserved family of DNA binding proteins that are | | | | Storage&Stability: |
| important in developmental regulation. The Kruppel zinc | | | | Store at 4 $^{\rm C}$ short term. Aliquot and store at -20 $^{\rm C}$ long |
| finger transcription factor was initially identified in Dro- | | | | term. Avoid freeze-thaw cycles. |
| sophila as a segmentation gene. Kruppel-like factors that | | | | Specificity: |
| have been characterized in mammals include EKLF, | | | clude EKLF, | EKLF (W318) polyclonal antibody detects endogenous |
| LKLF and | KLF and GKLF. EKLF is expressed principally in rythroid tissues, and LKLF expression is limited to the | | | levels of EKLF protein. |
| erythroid ti | | | | DATA: |

shown to be expressed during growth arrest.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

lung. GKLF is found predominantly in gut and has been

Molecular Weight:

~ 38 kDa

Swiss-Prot:

Q13351

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).



Western blot analysis of Rat liver, using EKLF pAb at 1:1000 dilution.

Note:

For research use only, not for use in diagnostic procedure.

| Bioworld Technology, Inc. | | | | |
|---------------------------|---|--|--|--|
| Add: | 1660 South Highway 100, Suite 500 St. Louis Park, | | | |
| | MN 55416,USA. | | | |
| Email: | info@bioworlde.com | | | |
| Tel: | 6123263284 | | | |
| Fax: | 6122933841 | | | |