

EphB6 (D881) polyclonal antibody

Catalog: BS2450

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Ephrin receptors and their ligands, the ephrins, mediate numerous developmental processes, particularly in the nervous system. Based on their structures and sequence relationships, ephrins are divided into the ephrin-A (EF-NA) class, which are anchored to the membrane by a glycosylphosphatidylinositol linkage, and the ephrin-B (EFNB) class, which are transmembrane proteins. The Eph family of receptors are divided into 2 groups based on the similarity of their extracellular domain sequences and their affinities for binding ephrin-A and ephrin-B ligands. Ephrin receptors make up the largest subgroup of the receptor tyrosine kinase (RTK) family. The ephrin receptor encoded by this gene lacks the kinase activity of most receptor tyrosine kinases and binds to ephrin-B ligands.

Product:

1 mg/ml in Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

Molecular Weight:

~ 119 kDa

Swiss-Prot:

015197

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200

IF: 1:50~1:200

Storage&Stability:

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at -20 $^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.

Specificity:

Bioworld Technology, Inc.

 Add:
 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.


 Email:
 info@bioworlde.com

 Tel:
 6123263284

 Fax:
 6122933841

EphB6 (D881) polyclonal antibody detects endogenous levels of EphB6 protein.

DATA:




Western blot (WB) analysis of EphB6 (D881) polyclonal antibody at 1:500 dilution

Lane1:Hela whole cell lysate

Lane2:NIH-3T3 whole cell lysate

Lane3:PC12 whole cell lysate

Lane4:H9C2 whole cell lkysate



Immunohistochemistry (IHC) analyzes of EphB6 (D881) pAb in paraffin-embedded human breast carcinoma tissue at 1:50.showing secreted and cell membrane staining. Negative control (the right)Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld technology, co. Ltd.

 Add:
 No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

 Email:
 info@biogot.com

 Tel:
 0086-025-68037686

 Fax:
 0086-025-68035151