

**DOR-1 (F345) polyclonal antibody**

Catalog: BS3179

Host: Rabbit

Reactivity: Human

BackGround:

The opioid receptors are G-protein coupled, seven-transmembrane domain receptors for enkephalins, endorphins, and dynorphins. Three different opioid receptor subtypes (kappa, delta, and mu) were first identified by their different selectivities for various naturally occurring alkaloid opioid ligands, and subsequently confirmed by molecular cloning. The amino acid sequences of the opioid receptor subtypes are ~70% homologous, and are similar to somatostatin receptors (SSTRs) showing ~40% homology with SSTR1. G-protein binding is thought to occur at the third intracellular loop of the opioid receptors, which is also the location of consensus sequences for phosphorylation of the receptor. Interestingly, the genes encoding the specific receptor subtypes are found on different chromosomes in both the human and mouse genomes.

Product:

1 mg/ml in Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

Molecular Weight:

~ 35, 50 kDa

Swiss-Prot:

P41143

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

IHC: 1:50~1:200

Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

DOR-1 (F345) polyclonal antibody detects endogenous levels of DOR-1 protein.

DATA:**Note:**

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151