

ACTR-IIA (H59) polyclonal antibody

Catalog: BS3669

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Members of the transforming growth factor β superfamily bind to a pair of transmembrane proteins, known as receptor types I and II, which contain serine/threonine kinases and associate to form a signaling complex. Activin has been shown to bind a heteromeric noncovalent complex, which consists of a type I receptor, ACTR-IA (also designated ACVRI and ALK-2) or ACTR-IB (also designated ALK-4 and SKR2), and a type II receptor, ACTR-IIA (ACVR2A) or ACTR-IIB (ACVR2B). Both receptor types are highly expressed in brain. The activin receptor family members are thought to mediate distinct effects on gene expression, cell differentiation and morphogenesis in a dose dependent manner.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 58 kDa

Swiss-Prot:

P27037

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

ACTR-IIA (H59) polyclonal antibody detects endogenous levels of ACTR-IIA protein.

DATA:



Western blot (WB) analysis of ACTR-IIA (H59) pAb at 1:500 dilution

Lane1: AML-12 whole cell lysate(40ug)

Lane2: C6 whole cell lysate(40ug)

Lane3: HEK293T whole cell lysate(40ug)

Lane4: A549 whole cell lysate(40ug)

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogol.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151