

2 3

180

130

95

eEF2K (phospho-S366) polyclonal antibody

Catalog: **BS4060** Host: Rabbit Reactivity: Human

p-eEF2K (\$366)

BackGround:

eEF2K is 105 kDa protein that is detected in skeletal muscle extracts and is phosphorylated rapidly by SAPK4, but poorly by p38, p38y, JNK or ERK 2. SAPK4 phosphorylates eEF2K at Ser 359 and Ser 396 in vitro, causing its inactivation. The phosphorylation of eEF2K at Ser 359 is also induced by insulin-like growth factor-1. Ser 359 is in close proximity to Ser 366 and the Ser 366 residue also becomes phosphorylated in response to growth factors. eEF2K is phosphorylated by p70 S6 kinase at Ser 366 and this results in the inactivation of eEF2K, especially at low (micromolar) calcium concentrations.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 105 kDa

Swiss-Prot:

O00418

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200

IP: 1:10~1:100

Storage&Stability:

Store at 4 $^{\circ}$ C short term. Aliquot and store at -20 $^{\circ}$ C long term. Avoid freeze-thaw cycles.

Specificity:

p-eEF2K (S366) polyclonal antibody detects endogenous levels of eEF2K protein when phosphorylated at Ser366.

DATA:

Add:

1660 South Highway 100, Suite 500 St. Louis Park, Add: P. R. China.

Fax:

MN 55416,USA. Email: info@bioworlde.com Tel: 6123263284

Bioworld Technology, Inc.

6122933841 Fax:

Bioworld technology, co. Ltd. No 9, weidi road Qixia District Nanjing, 210046, **Email:** info@biogot.com Tel: 0086-025-68037686 0086-025-68035151





Immunohistochemistry (IHC) analyzes of p-eEF2K (S366) pAb in paraffin-embedded human breast carcinoma tissue at 1:100.



Immunoprecipitation of HEK293T cell lysate using p-eEF2K (S366) pAb (Sepharose Bead Conjugate) #BD0048(lane 2 and lane 3) and Nonspecific IgG Control (Sepharose Bead Conjugate)#BD0048 (lane 4) .Lane 1 is 30% input. The western blot was probed using p-eEF2K (S366). " \uparrow " (supernatant) ; " \downarrow (deposition)

Note:

For research use only, not for use in diagnostic procedure.