

IκB-α (phospho-Y42) polyclonal antibody

Catalog: BS4736

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Activation of NFκB requires that IκB be phosphorylated on specific serine residues, which results in targeted degradation of IκB. IκB kinase α (IKKα), previously designated CHUK, interacts with IκB-α and specifically phosphorylates IκB-α on the sites that trigger its degradation Serines 32 and 36. IKKα appears to be critical for NFκB activation in response to proinflammatory cytokines. Phosphorylation of IκB by IKKα is stimulated by the NFκB inducing kinase (NIK), which itself is a central regulator for NFκB activation in response to TNF and IL-1. The functional IKK complex contains three subunits, IKKα, IKKβ and IKKγ, and each appear to make essential contributions to IκB phosphorylation.

Product:

1 mg/ml in Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

Molecular Weight:

~ 36 kDa

Swiss-Prot:

P25963

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

p-IκB-α (Y42) polyclonal antibody detects endogenous levels of IκB-α protein only when phosphorylated at Tyr42.

DATA:



Western blot (WB) analysis of p-IκB-α (Y42) pAb at 1:500 dilution


Lane1: The Brain tissue lysate of Mouse(40ug)

Lane2: SGC7901 whole cell lysate(40ug)

Lane3: HCT116 whole cell lysate(40ug)

Lane4: MCF-7 whole cell lysate(40ug)

Lane5: Beas-2B whole cell lysate(40ug)



Immunohistochemistry (IHC) analyzes of p-IκB-α (Y42) pAb in paraffin-embedded human breast carcinoma tissue.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151