

STAM2 (phospho-Y192) polyclonal antibody

Catalog: BS4876

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

STAM contains an SH3 (Src homology 3) motif as well as an immunoreceptor tyrosine-based activation (ITAM) motif, both of which appear to be required for c-Myc induction in response to IL-2 and GM-CSF. STAM associates with JAK3 and JAK2 via its ITAM region, and it is tyrosine phosphorylated by JAK3 and JAK2 after stimulation with IL-2 and GM-CSF, respectively. STAM2, also known as Hbp, is a protein that is highly related to STAM. Similar to STAM, STAM2 functions downstream of JAK kinases and can be phosphorylated in response to cytokines. Due to alternative splicing events, two isoforms of STAM2 exist, namely STAM2A and STAM2B.

Product:

1 mg/ml in Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

Molecular Weight:

~ 58 kDa

Swiss-Prot:

O75886

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

p-STAM2 (Y192) polyclonal antibody detects endogenous levels of STAM2 protein only when phosphorylated at Tyr192.

DATA:



Western blot (WB) analysis of p-STAM2 (Y192) polyclonal antibody at 1:500 dilution

Lane1: HEK293T cell treated with EGF(0.1ng/ML, 30min)

Lane2: sp2/0 cell treated with EGF(0.1ng/ML, 30min)

Lane3: Rat spleen tissue lysate

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151