

DNA pol δ 3 polyclonal antibody

Catalog: BS5699

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

DNA replication, recombination and repair, all of which are necessary for genomic stability, require the presence of exonucleases. In DNA replication, exonucleases are involved in the processing of Okazaki fragments, whereas in DNA repair, they function to excise damaged DNA fragments and correct recombinational mismatches. These exonucleases include the family of DNA polymerases, namely DNA pol α , β , δ and ϵ . DNA pol δ and DNA pol ϵ are multisubunit enzymes, with DNA pol δ consisting of two subunits which interact with the sliding DNA clamp protein PCNA. DNA pol δ 3, also known as POLD3 or p66, is a 466 amino acid nuclear protein that is required for the proper function of DNA pol δ .

Product:

1mg/ml in PBS with 0.1% Sodium Azide, 50% Glycerol.

Molecular Weight:

~ 51 kDa

Swiss-Prot:

Q15054

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

ICC: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

DNA pol δ 3 polyclonal antibody detects endogenous levels of DNA pol δ 3 protein.

DATA:



Western blot (WB) analysis of DNA pol δ 3 polyclonal antibody at 1:500 dilution

Lane1:Hela cell lysate

Lane2:sp2/0 cell lysate

Lane3:H9C2 cell lysate

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogol.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151