

IAP polyclonal antibody

Catalog: BS60372

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

The intestinal alkaline phosphatase gene (ALPI) encodes a digestive brush-border enzyme, IAP (also designated ALP), which is highly upregulated during small intestinal epithelial cell differentiation. IAP, an enterocyte differentiation marker that functions to limit fat absorption, has been implicated in trans-cellular transport of chylomicrons and in chylomicron formation. At high pH, IAP removes phosphate groups from proteins and from the 5' end of DNA and RNA. Most mammals have four different IAP isozymes: placental, placental-like, intestinal and non tissue-specific. Non tissue-specific isozymes are found in liver, kidney and bone. Tissues with particularly high concentrations of IAP include the liver, bile ducts, placenta and bone.

Product:

1mg/ml in PBS with 0.1% Sodium Azide, 50% Glycerol.

Molecular Weight:

~ 57 kDa

Swiss-Prot:

P09923

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

IAP polyclonal antibody detects endogenous levels of IAP protein.

DATA:



Western blot (WB) analysis of IAP polyclonal antibody at 1:500 dilution

Lane1:HEK293T cell lysate

Lane2:Raw264.7 cell lysate

Lane3:PC12 cell lysate

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151