

DFFB polyclonal antibody

Catalog: BS60830

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

The Ced/ICE or caspase family of cysteine proteases plays a pivotal role in mediating apoptosis through the proteolysis of specific targets. Among the targets are poly(ADP-ribose) polymerase (PARP), Gelsolin, ICAD (inhibitor of CAD, also designated DFF-45) and the nuclear lamins. CAD (caspase-activated deoxyribonuclease), also designated CPAN (caspase-activated nuclease) and DFF40, is a DNase that is responsible for DNA degradation during apoptosis. Caspase-3 acts to dissociate CAD from ICAD, allowing CAD to enter the nucleus and degrade DNA.

Product:

1 mg/ml in Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.3.

Molecular Weight:

~ 39 kDa

Swiss-Prot:

O76075

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:


WB: 1:500~1:1000

Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

DFFB polyclonal antibody detects endogenous levels of DFFB protein.

DATA:

Western blot (WB) analysis of DFFB polyclonal antibody at 1:500 dilution

Lane1:HEK293T whole cell lysate

Lane2:MCF-7 whole cell lysate

Lane3:H9C2 whole cell lysate

Lane4:Raw264.7 whole cell lysate

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogol.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151