

RIP polyclonal antibody

Catalog: BS91172

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

In contrast to growth factors which promote cell proliferation, FAS ligand (FAS-L) and the tumor necrosis factors (TNFs) rapidly induce apoptosis. Cellular response to FAS-L and TNF is mediated by structurally related receptors containing a conserved "death domain" and belonging to the TNF receptor superfamily. TRADD, FADD and RIP are FAS/TNF-R1 interacting proteins that contain a death domain homologous region (DDH). TRADD (TNF-R1-associated death domain) and FADD (FAS-associated death domain) associate with the death domains of both FAS and TNF-R1 via their DDH regions. Overexpression of TRADD leads to NF κ B activation and apoptosis in the absence of TNF. Overexpression of FADD causes apoptosis, which can be blocked by the cow pox protein CrmA, suggesting that FADD lies upstream of ICE and possibly other serine proteases. The receptor interacting protein, RIP, associates with FAS exclusively via its DDH and this association is abrogated in *lpr* mutants. Unlike TRADD and FADD, RIP contains a putative amino terminal kinase domain.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 76-90 kDa

Swiss-Prot:

Q13546(Human)

Purification&Purity:

ProA affinity purified

Applications:

WB:1:1,000-1:5,000

FC:1:50-1:100

Storage&Stability:

Store at +4 °C after thawing. Aliquot store at -20 °C or -80 °C. Avoid repeated freeze / thaw cycles.

Specificity:

RIP polyclonal antibody detects endogenous levels of RIP protein.

DATA:



Western blot (WB) analysis of RIP pAb at 1:1000 dilution

Lane1:C6 whole cell lysate(40ug)


Lane2:3T3-L1 whole cell lysate(40ug)

Lane3:The Brain tissue lysate of Mouse(40ug)

Lane4:The Stomach tissue lysate of Rat(40ug)

Lane5:DLD whole cell lysate(40ug)

Lane6:SGC7901 whole cell lysate(40ug)



Flow cytometric analysis of 293 cells with RIP antibody at 1/50 dilution

(red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151



PRODUCT DATA SHEET

Bioworld Biotech Co., Ltd

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,
MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,
P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151