

HC-II (D68) polyclonal antibody

Catalog: BS9125

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Heparin cofactor II (HCII) is a glycoprotein in human plasma which rapidly inactivates Thrombin in the presence of dermatan sulfate. Inhibition occurs by formation of a stable equimolar complex between HCII and Thrombin. Certain clinical conditions, such as hepatic failure, disseminated intravascular coagulation, thalassemia and sickle cell anemia, display reduced levels of HCII. However, during pregnancy, physiological levels of HCII expression are elevated. HCII may regulate coagulation and may participate in processes such as inflammation, atherosclerosis and wound repair. HCII is widely distributed among vertebrates and may have a common function in birds, amphibians and mammals.

Product:

1 mg/ml in Phosphate buffered saline (PBS) with 0.05% sodium azide, approx. pH 7.2.

Molecular Weight:

~ 57 kDa

Swiss-Prot:

P05546

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

IHC: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

HC-II (D68) polyclonal antibody detects endogenous levels of HC-II protein.

DATA:



Immunohistochemistry (IHC) analyzes of HC-II (D68) pAb in paraffin-embedded human breast carcinoma tissue at 1:100.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151