

CYP17A1 monoclonal antibody

Catalog: MB0132

Host: Mouse

Reactivity: Human

BackGround:

CYP17A (17 α -hydroxylase/17,20-lyase) is important for the conversion of pregnenolone and progesterone to dehydroepiandrosterone (DHEA) and androstenedione. In this process, it catalyzes both the 17 α -hydroxylation and the 17,20-lyase reaction. CYP17A1 is crucial during sexual development, both during fetal development and during puberty, and is intracellularly regulated by cAMP levels. Defects in the CYP17A1 gene, which encodes for the protein, may cause adrenal hyperplasia type V (AH-V) which is characterized by hypokalemia and hypertension. Male patients affected by AH-V do not undergo normal sexual differentiation and develop female external genitalia and do not undergo pubertal development.

Product:

Purified mouse monoclonal in buffer containing 0.1M Tris-Glycine (pH 7.4, 150 mM NaCl) with 0.2% sodium azide, 50% glycerol

Molecular Weight:

Predicted band size: 57KDa

Observed band size: 57KDa

Swiss-Prot:

P05093

Purification&Purity:

The antibody was affinity-purified from mouse ascites by

affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:1000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

This antibody detects endogenous levels of CYP17A1 and does not cross-react with related proteins.

DATA:



Western blot detection of CYP17A1 antibody in HeLa, HepG2 and A549 cell lysates using CYP17A1 antibody (1:1000 diluted).

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogol.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151