

CD336 Recombinant Protein

Catalog: NCP0296

Host: E.coli

Tag: His-tag

BackGround:

Natural killer (NK) cells direct cytotoxicity against tumor or virally infected cells. NK cell-mediated cytotoxicity is stimulated by several activating receptors associated with the signaling adapter DNAX activation 12/killer cell-activating receptor-associated protein (DAP12). NKp44 is a natural cytotoxicity receptor that is expressed on IL-2-activated human NK cells and may contribute to the increased efficiency of NK cells to mediate tumor cell lysis. NKp44 is composed of one Ig-like extracellular domain, a transmembrane segment and a cytoplasmic domain. Prolactin upregulates and cortisol downregulates the surface expression of NKp44 at the transcriptional level. A cellular ligand for NKp44 (NKp44L) is expressed during HIV-1 infection and is correlated with the progression of CD4+ T cell depletion and an increase of viral load. This implicates NKp44 as a therapeutic agent that may aid in the progress towards a vaccine for HIV-1 infection.

Product:

PBS, 4M Urea, PH7.4

Molecular Weight:

~19kDa

Swiss-Prot:

O95944

Purification&Purity:

Transferred into competent cells and the supernatant was purified by NI column affinity chromatography and the purity is > 85% (by SDS-PAGE).

Restriction Sites:

NdeI-XhoI


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Expression Vector:

pet-22b(+)

DATA:



SDS-PAGE Result

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,
MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,
P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151