PRODUCT DATA SHEET



Bioworld Technology, Inc.

Recombinant Human Thrombopoietin (rHuTPO)

Catalog Number: PR1120 Source:CHO Quantity:2µg/10µg/1.0mg

Description

Thrombopoietin (Tpo), the ligand for the receptor encoded by the c-Mpl proto-oncogene, is a key regulator of megakaryocytopoiesis and thrombopoiesis in vitro and in vivo. The cD-NAs for Tpo have recently been cloned from canine, murine and human sources. The proteins from these three species are highly conserved, exhibiting from 69-75% sequence identity at the amino acid level. Human Tpo cDNA encodes a 353 amino acid residue protein with a 21 amino acid residue signal peptide that is cleaved to yield the 332 amino acid residue mature protein. Two distinct domains, separated by a pair of arginine residues that may be a proteolytic cleavage site, have been identified in Tpo: the amino terminal region exhibiting sequence homology to erythropoietin and the carboxy terminal region containing multiple potential N-linked glycosylation sites. Recombinant Tpo has now been shown to stimulate the maturation, as well as the proliferation, of megakaryocytes and may have important therapeutic applications for the treatment of various clinical conditions associated with thrombocytopenia.

Molecular Weight:

Approximately 80 kDa, consisting of a 332 amino acid residue with a predicted molecular mass of approximately 35 kDa. As a result of glycosylation, the recombinant protein migrates with an apparent molecular mass of 80±10 kDa in SDS-PAGE.

Purity:

>98% by SDS-PAGE and HPLC analyses.

Biological Activity:

Fully biologically active when compared to standard. The ED50 as determined by the dose-dependant stimulation of Mo7e cells is typically 1-3 ng/ml, corresponding to a Specific Activity of \Box 1 x 10 6 IU/mg.

Physical Appearance:

Sterile Filtered White lyophilized (freeze-dried) powder.

Formulation:

Lyophilized from a 0.2mm filtered concentrated solution in PBS, pH 7.4.

AA Sequence:

SPAPPACDLRVLSKLLRDSHVLHSRLS
QCPEVHPLPTPVLLPAVDFSLGEWKTQ
MEETKAQDILGAVTLLLEGVMAARGQ
LGPTCLSSLLGQLSGQVRLLLGALQSL
LGTQLPPQGRTTAHKDPNAIFLSFQHL
LRGKVRFLMLVGGSTLCVRRAPPTTAV
PSRTSLVLTLNELPNRTSGLLETNFTAS
ARTTGSGLLKWQQGFRAKIPGLLNQTS
RSLDQIPGYLNRIHELLNGTRGLFPGPS
RRTLGAPDISSGTSDTGSLPPNLQPGYS
PSPTHPPTGQYTLFPLPPTLPTPVVQLH
PLLPDPSAPTPTPTSPLLNTSYTHSQNL
SQEG

Endotoxin:

Less than 1EU/mg of rHuTPO a as determined by LAL method.

Reconstitution:

Tel: 0086-025-86371664

We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in

MADE IN CHINA

Bioworld technology, co, Ltd.

No 9, weidi road Qixia District Nanjing, 210046, P, R.China.

Email: info@biogot.com Fax:0086-025-86213570

1660 South Highway 100, Suite 500 St. Louis Park, MN

Bioworld Technology, Inc.

PRODUCT DATA SHEET

sterile distilled water or aqueous buffer containing 0.1% BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at <-20°C. Further dilutions should be made in appropriate buffered solutions.

Storage:

This lyophilized preparation is stable at 2-8°C, but should be kept at -20°C for long term storage, preferably desiccated.

Bioworld Technology, Inc.

1660 South Highway 100, Suite 500 St. Louis Park, MN

55416,USA.

Tel: 6123263284

Upon reconstitution, the preparation is stable for up to one week at 2-8°C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20°C to -70°C. Avoid repeated freeze/thaw cycles.

Usage:

This material is offered by USA Bioworld biotech for research, laboratory or further evaluation purposes. NOT FOR HUMAN USE. Made in China

Email: info@bioworlde.com Fax: 6122933841 Tel: 0086-025-86371664