

ABHD8 polyclonal antibody

Catalog: BS30855

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Protein ABHD8: Negatively regulates NLRP3-driven inflammation. Promotes NLRP3 degradation through the chaperone-mediated autophagy (CMA) pathway, hence attenuating inflammasome activation and IL1B secretion. Acts by recruiting palmitoyltransferase ZDHHC12 to NLRP3, facilitating NLRP3 palmitoylation and subsequent degradation.

Product:

1mg/ml in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

Molecular Weight:

~47.3 kDa

Swiss-Prot:

Q96I13

Purification&Purity:

The antibody was purified by immunogen affinity chromatography.

Applications:

WB (1/500 - 1/1000), IHC (1/50 - 1/100)


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Recognizes endogenous levels of ABHD8 protein.

DATA:



Western blot analysis of SP1 (Phospho-T453) expression in H1688 (A) whole cell lysates.



Immunohistochemical analysis of SP1 (Phospho-T453) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX. Immunofluorescent analysis of SP1 (Phospho-T453) staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151