

AE4 polyclonal antibody

Catalog: BS30859

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Anion exchange protein 4: Electroneutral $\text{Cl}(-)/\text{HCO}_3(-)$ antiporter that favors chloride ion entry and efflux of hydrogencarbonate and sodium ion across the basolateral membrane and may participate in salivary secretion. Also mediates $\text{Cl}(-)/\text{HCO}_3(-)$ exchange activity in the presence of $\text{K}(+)$ as well as $\text{Cs}(+)$, $\text{Li}(+)$, and $\text{Rb}(+)$ (By similarity). Does not contribute to $\text{Cl}(-)/\text{HCO}_3(-)$ exchanger in the apical membrane of the upper villous epithelium (By similarity)

Product:

1mg/ml in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

Molecular Weight:

~108 kDa

Swiss-Prot:

Q96Q91

Purification&Purity:

The antibody was purified by immunogen affinity chromatography.

Applications:

WB (1/500 - 1/1000)


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Recognizes endogenous levels of AE4 protein.

DATA:



Immunohistochemical analysis of Cytokeratin 19 staining in human kidney formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Immunohistochemical analysis of Cytokeratin 19 staining in human appendix formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151