

PTCHD4 polyclonal antibody

Catalog: BS31548

Host: Rabbit

Reactivity: Human

BackGround:

Patched domain-containing protein 4: Could act as a repressor of canonical hedgehog signaling by antagonizing the effects of SMO, as suggested by down-regulation of hedgehog target genes, including GLI1, PTCH1, and PTCH2 in PTCHD4-expressing cells

Product:

1mg/ml in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

Molecular Weight:

~96.4 kDa

Swiss-Prot:

Q6ZW05

Purification&Purity:

The antibody was purified by immunogen affinity chromatography.

Applications:

WB (1/500 - 1/1000), IHC (1/50 - 1/200)


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.


Specificity:

Recognizes endogenous levels of PTCHD4 protein.

DATA:



Western blot analysis of Factor XIIIa expression in mouse testis (A), rat testis (B), K562 (C), EC9706 (D), MCF7 (E), HCT116 (F) whole cell lysates.



Immunohistochemical analysis of Factor XIIIa staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151