



## PRODUCT DATA SHEET

Bioworld Technology, Inc.

### TTLL4 polyclonal antibody

Catalog: BS31670

Host: Rabbit

Reactivity: Human

#### BackGround:

Tubulin monoglutamylase TTLL4: Monoglutamylase which modifies both tubulin and non-tubulin proteins, adding a single glutamate on the gamma-carboxyl group of specific glutamate residues of target proteins. Involved in the side-chain initiation step of the polyglutamylation reaction but not in the elongation step. Preferentially modifies beta-tail tubulin over the alpha-tubulin. Monoglutamylates nucleosome assembly proteins NAP1L1 and NAP1L4. Monoglutamylates nucleotidyl-transferase CGAS, leading to inhibition of CGAS catalytic activity, thereby preventing antiviral defense function. Involved in KLF4 glutamylation which impedes its ubiquitination, thereby leading to somatic cell reprogramming, pluripotency maintenance and embryogenesis

#### Product:

1mg/ml in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

#### Molecular Weight:

~133 kDa

#### Swiss-Prot:

Q14679

#### Purification&Purity:

The antibody was purified by immunogen affinity chromatography.

#### Applications:

WB (1/500 - 1/1000)


#### Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

#### Specificity:

Recognizes endogenous levels of TTLL4 protein.

#### DATA:



Western blot analysis of TIP47 expression in HeLa (A), K562 (B), Daudi (C) whole cell lysates.

Flow cytometric analysis of HUVEC cells using Anti-TIP47 Antibody.

The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody at 37 °C for 60 min. The secondary antibody Goat Anti-Mouse IgG (H&L) - AF488 was incubated at 37 °C for 40 min. Isotype control antibody (blue line) was used under the same condition.

#### Note:

For research use only, not for use in diagnostic procedure.

#### Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: [info@bioworld.com](mailto:info@bioworld.com)

Tel: 6123263284

Fax: 6122933841

#### Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: [info@biogot.com](mailto:info@biogot.com)

Tel: 0086-025-68037686

Fax: 0086-025-68035151