

RIC3 polyclonal antibody

Catalog: BS31687

Host: Rabbit

Reactivity: Human

BackGround:

Protein RIC-3: Molecular chaperone which facilitates proper subunit assembly and surface trafficking of alpha-7 (CHRNA7) and alpha-8 (CHRNA8) nicotinic acetylcholine receptors. May also promote functional expression of homomeric serotonergic 5-HT3 receptors, and of heteromeric acetylcholine receptors alpha-3/beta-2, alpha-3/beta-4, alpha-4/beta-2 and alpha-4/beta-4

Product:

1mg/ml in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

Molecular Weight:

~41.1 kDa

Swiss-Prot:

Q7Z5B4

Purification&Purity:

The antibody was purified by immunogen affinity chromatography.

Applications:

WB (1/500 - 1/1000), IHC (1/50 - 1/200)


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.


Specificity:

Recognizes endogenous levels of RIC3 protein.

DATA:



Western blot analysis of Complement C3 expression in HepG2 (A) whole cell lysates.



Immunohistochemical analysis of Complement C3 staining in human liver cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Immunofluorescent analysis of Complement C3 staining in rat liver. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AF594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151