

DHRS7C polyclonal antibody

Catalog: BS31807

Host: Rabbit

Reactivity: Human

BackGround:

Dehydrogenase/reductase SDR family member 7C: NADH-dependent oxidoreductase which catalyzes the oxidation of all-trans-retinol to all-trans-retinal. Plays a role in the regulation of cardiac and skeletal muscle metabolic functions. Maintains Ca(2+) intracellular homeostasis by repressing Ca(2+) release from the sarcoplasmic reticulum (SR) in myotubes, possibly through local alterations in NAD/NADH or retinol/retinal. Also plays a role in Ca(2+) homeostasis by controlling Ca(2+) overload in the cytosol and the SR in myotubes. Involved in glucose uptake into skeletal muscles and muscle performance by activating PI3K and mTORC2-mediated AKT1 phosphorylation signaling pathways, possibly through the action of its downstream catalytic product all-trans-retinoic acid

Product:

1mg/ml in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

Molecular Weight:

~34.9 kDa

Swiss-Prot:

A6NNS2

Purification&Purity:

The antibody was purified by immunogen affinity chromatography.

Applications:

WB (1/500 - 1/1000), IHC (1/50 - 1/200)



Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Recognizes endogenous levels of DHRS7C protein.

DATA:



Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151