

WDR20 polyclonal antibody

Catalog: BS32004

Host: Rabbit

Reactivity: Human

BackGround:

WD repeat-containing protein 20: Regulator of deubiquitinating complexes. Activates deubiquitinating activity of complexes containing USP12. Anchors at the base of the ubiquitin-contacting loop of USP12 and remotely modulates the catalytic center of the enzyme. Regulates shuttling of the USP12 deubiquitinase complex between the plasma membrane, cytoplasm and nucleus

Product:

1mg/ml in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

Molecular Weight:

~62.9 kDa

Swiss-Prot:

Q8TBZ3

Purification&Purity:

The antibody was purified by immunogen affinity chromatography.

Applications:

WB (1/500 - 1/1000)


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Recognizes endogenous levels of WDR20 protein.

DATA:



Western blot analysis of Endothelin 1 expression in HEK293T (A), LOVO (B), H1792 (C), mouse testis (D) whole cell lysates.

Immunohistochemical analysis of Endothelin 1 staining in human colorectal cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151