

RBM24 polyclonal antibody

Catalog: BS32388

Host: Rabbit

Reactivity: Human, Mouse

BackGround:

RNA-binding protein 24: Multifunctional RNA-binding protein involved in the regulation of pre-mRNA splicing, mRNA stability and mRNA translation important for cell fate decision and differentiation. Plays a major role in pre-mRNA alternative splicing regulation. Mediates preferentially muscle-specific exon inclusion in numerous mRNAs important for striated cardiac and skeletal muscle cell differentiation. Binds to intronic splicing enhancer (ISE) composed of stretches of GU-rich motifs localized in flanking intron of exon that will be included by alternative splicing (By similarity). Involved in embryonic stem cell (ESC) transition to cardiac cell differentiation by promoting pre-mRNA alternative splicing events of several pluripotency and/or differentiation genes. Plays a role in the regulation of mRNA stability. Binds to 3'-untranslated region (UTR) AU-rich elements in target transcripts, such as CDKN...

Product:

1mg/ml in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

Molecular Weight:

~24.8 kDa

Swiss-Prot:

Q9BX46

Purification&Purity:

The antibody was purified by immunogen affinity chromatography.

Applications:

WB (1/500 - 1/1000), IHC (1/50 - 1/200), FC (1/10 - 1/50)


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.



Specificity:

Recognizes endogenous levels of RBM24 protein.

DATA:



Western blot analysis of SPINK8 expression in K562 (A) whole cell lysates.



Flow cytometric analysis of K562 cells using Anti-SPINK8 Antibody. The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody at 37 °C for 60 min. The secondary antibody Goat Anti-Rabbit IgG (H&L) - AF488 was incubated at 37 °C for 40 min. Isotype control antibody (blue line) was used under the same condition.

Note:

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151



PRODUCT DATA SHEET

Bioworld Technology, Inc.

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,
MN 55416, USA.

Email: info@bioworldde.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,
P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151