



PRODUCT DATA SHEET

Bioworld Technology, Inc.

CYB561 polyclonal antibody

Catalog: BS32855

Host: Rabbit

Reactivity: Human, Mouse

BackGround:

Transmembrane ascorbate-dependent reductase CYB561: Transmembrane reductase that probably uses ascorbate as an electron donor to reduce Fe(3+) into Fe(2+) and could play a role in iron transport. It is also able to reduce extracellular monodehydro-L-ascorbate and could be involved in the regeneration and homeostasis within secretory vesicles of ascorbate which in turn provides reducing equivalents needed to support the activity of intravesicular enzymes

Product:

1mg/ml in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

Molecular Weight:

~27.6 kDa

Swiss-Prot:

P49447

Purification&Purity:

The antibody was purified by immunogen affinity chromatography.

Applications:

WB (1/500 - 1/1000)

Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Recognizes endogenous levels of CYB561 protein

DATA:

Western blot analysis of LZK expression in EC9706 (A), SGC7901 (B), Panc1 (C), AML12 (D), rat brain (E), HepG2 (F), LO2 (G) whole cell lysates.

Immunohistochemical analysis of LZK staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Immunofluorescent analysis of LZK staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151