



## PRODUCT DATA SHEET

Bioworld Technology, Inc.

### SOGA3 polyclonal antibody

Catalog: BS33404

Host: Rabbit

Reactivity: Human, Mouse

#### BackGround:

Microtubule cross-linking factor 3. This protein plays important roles in cellular processes, and its expression and regulation are of interest in biomedical research.

#### Product:

1mg/ml in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

#### Molecular Weight:

~103 kDa

#### Swiss-Prot:

Q5TF21

#### Purification&Purity:

The antibody was purified by immunogen affinity chromatography.

#### Applications:

WB (1/500 - 1/1000)

#### Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

#### Specificity:

Recognizes endogenous levels of SOGA3 protein

#### DATA:

Western blot analysis of SRP72 expression in Hela (A), K562 (B) whole cell lysates.

Immunohistochemical analysis of SRP72 staining in human pancreas formalin fixed paraffin embedded tissue section. The section was

pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Immunofluorescent analysis of Anti-SRP72 staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Flow cytometric analysis of Hela cells using Anti-SRP72 Antibody. The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody at 37 °C for 60 min. The secondary antibody Goat Anti-Rabbit IgG (H&L) - AF488 was incubated at 37 °C for 40 min. Isotype control antibody (blue line) was used under the same condition.

#### Note:

For research use only, not for use in diagnostic procedure.

#### Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: [info@bioworld.com](mailto:info@bioworld.com)

Tel: 6123263284

Fax: 6122933841

#### Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: [info@biogot.com](mailto:info@biogot.com)

Tel: 0086-025-68037686

Fax: 0086-025-68035151