



CIPC polyclonal antibody

Catalog: BS33791

Host: Rabbit

Reactivity: Human, Mouse

BackGround:

CLOCK-interacting pacemaker: Transcriptional repressor which may act as a negative-feedback regulator of CLOCK-BMAL1 transcriptional activity in the circadian-clock mechanism. May stimulate BMAL1-dependent phosphorylation of CLOCK. However, the physiological relevance of these observations is unsure, since experiments in an animal model showed that CIPC is not critically required for basic circadian clock

Product:

1mg/ml in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

Molecular Weight:

~42.7 kDa

Swiss-Prot:

Q9C0C6

Purification&Purity:

The antibody was purified by immunogen affinity chromatography.

Applications:

WB (1/500 - 1/1000)

Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Recognizes endogenous levels of CIPC protein

DATA:

Western blot analysis of CGB3 expression in HEK293T (A) whole cell lysates.

Immunohistochemical analysis of CGB3 staining in human stomach formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151