



PRODUCT DATA SHEET

Bioworld Technology, Inc.

SPEF2 polyclonal antibody

Catalog: BS34229

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Sperm flagella and cilia-associated protein 2: Required for correct axoneme development in spermatozoa. Important for normal development of the manchette and sperm head morphology. Essential for male fertility. Plays a role in localization of the intraflagellar transport protein IFT20 to the manchette, suggesting function as an adapter for dynein-mediated protein transport during spermatogenesis. Plays a role in bone growth where it seems to be required for normal osteoblast differentiation (By similarity). Required for normal beat frequency of tracheal epithelial cilia and oviduct cilia (By similarity)

Product:

1mg/ml in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

Molecular Weight:

~210 kDa

Swiss-Prot:

Q9C093

Purification&Purity:

The antibody was purified by immunogen affinity chromatography.

Applications:

WB (1/500 - 1/1000)

Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Recognizes endogenous levels of SPEF2 protein

DATA:

Western blot analysis of HSP70 expression in THP1 (A) whole cell lysates.

Immunohistochemical analysis of HSP70 staining in human prostate formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Immunofluorescent analysis of HSP70 staining in THP1 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151