

**ATP6V1G1 polyclonal antibody**

Catalog: BS34509

Host: Rabbit

Reactivity: Human, Mouse

BackGround:

V-type proton ATPase subunit G 1: Subunit of the V1 complex of vacuolar(H⁺)-ATPase (V-ATPase), a multi-subunit enzyme composed of a peripheral complex (V1) that hydrolyzes ATP and a membrane integral complex (V0) that translocates protons . V-ATPase is responsible for acidifying and maintaining the pH of intracellular compartments and in some cell types, is targeted to the plasma membrane, where it promotes acidification of the extracellular environment . The V-ATPase complex also acts as an activator for mTORC1 on lysosomal membrane by promoting the guanine nucleotide exchange factor (GEF) of the Ragulator complex, thereby enabling mTORC1 recruitment . In aerobic conditions, involved in intracellular iron homeostasis, thus triggering the activity of Fe(2⁺) prolyl hydroxylase (PHD) enzymes, and leading to HIF1A hydroxylation and subsequent proteasomal degradation

Product:

1mg/ml in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

Molecular Weight:

~13.8 kDa

Swiss-Prot:

O75348

Purification&Purity:

The antibody was purified by immunogen affinity chromatography.

Applications:

WB (1/500 - 1/1000)

Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Recognizes endogenous levels of ATP6V1G1 protein

DATA:

Western blot analysis of DDX3X (Phospho-T322) expression in HeLa TNFa-treated (A) whole cell lysates.

Immunohistochemical analysis of DDX3X (Phospho-T322) staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Immunofluorescent analysis of DDX3X (Phospho-T322) staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151