



ZGLP1 polyclonal antibody

Catalog: BS34955

Host: Rabbit

Reactivity: Human, Mouse

BackGround:

GATA-type zinc finger protein 1: Transcriptional regulator that plays a key role in germ cell development. Determines the oogenic fate by activating key genes for the oogenic program and meiotic prophase entry. Acts downstream of bone morphogenetic protein (BMP) by regulating expression of genes required for the oogenic programs, which are repressed by Polycomb activities in sexually uncommitted germ cells. Regulates expression of STRA8, a central downstream effector for the meiotic program. Acts independently of retinoic acid (RA). In males, not required for germ-cell sex determination, but required to allow the spermatogonia to efficiently accomplish the meiotic prophase

Product:

1mg/ml in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

Molecular Weight:

~29.7 kDa

Swiss-Prot:

P0C6A0

Purification&Purity:

The antibody was purified by immunogen affinity chromatography.

Applications:

WB (1/500 - 1/1000)

Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Recognizes endogenous levels of ZGLP1 protein

DATA:

Western blot analysis of Aurora A expression in HEK293 (A), MCF7 (B), Hela (C) whole cell lysates.

Immunohistochemical analysis of Aurora A staining in human cervical cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Immunofluorescent analysis of Aurora A staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with an AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Flow cytometric analysis of HeLa cells using Anti-Aurora A Antibody (green) and negative control (red).

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151