



ZNF658 polyclonal antibody

Catalog: BS35256

Host: Rabbit

Reactivity: Human

BackGround:

Zinc finger protein 658: Mediates transcriptional repression in response to zinc. Represses several genes, including SLC30A5, SLC30A10 and CBWD1, by binding to the zinc transcriptional regulatory element (ZTRE) (5'-C[AC]C[TAG]CC[TC]-N(0-50)-[GA]G[ATC]G[TG]G-3') found in the promoter region. May play a role in the control of ribosome biogenesis, regulating predominantly rRNA levels, as well as those of several ribosomal proteins, thus coordinating this highly zinc-demanding process with the available zinc supply

Product:

1mg/ml in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

Molecular Weight:

~122 kDa

Swiss-Prot:

Q5TYW1

Purification&Purity:

The antibody was purified by immunogen affinity chromatography.

Applications:

WB (1/500 - 1/1000)

Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Recognizes endogenous levels of ZNF658 protein

DATA:

Western blot analysis of STK17B expression in MCF7 (A), K562 (B), HEK293T (C), mouse brain (D), rat brain (E) whole cell lysates.

Immunohistochemical analysis of STK17B staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Immunofluorescent analysis of STK17B staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark. Phalloidin - AF594 was used to stain Actin filaments (red). DAPI was used to stain the cell nuclei (blue).

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151