

**OTUD5 polyclonal antibody**

Catalog: BS35296

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

OTU domain-containing protein 5: Deubiquitinating enzyme that functions as a negative regulator of the innate immune system. Has peptidase activity towards 'Lys-48'- and 'Lys-63'-linked polyubiquitin chains. Can also cleave 'Lys-11'-linked ubiquitin chains (in vitro). Acts via TRAF3 deubiquitination and subsequent suppression of type I interferon (IFN) production. Controls neuroectodermal differentiation through cleaving 'Lys-48'-linked ubiquitin chains to counteract degradation of select chromatin regulators such as ARID1A, HDAC2 and HCF1. Acts as a positive regulator of mTORC1 and mTORC2 signaling following phosphorylation by MTOR: acts by mediating deubiquitination of BTRC, leading to its stability.

Product:

1mg/ml in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

Molecular Weight:

~60.6 kDa

Swiss-Prot:

Q96G74

Purification&Purity:

The antibody was purified by immunogen affinity chromatography.

Applications:

WB (1/500 - 1/1000)

Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Recognizes endogenous levels of OTUD5 protein

DATA:

Western blot analysis of Uromodulin expression in K562 (A), HEK293T (B) whole cell lysates.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151