



PRODUCT DATA SHEET

Bioworld Technology, Inc.

MGARP polyclonal antibody

Catalog: BS35563

Host: Rabbit

Reactivity: Human, Mouse

BackGround:

Protein MGARP: Plays a role in the trafficking of mitochondria along microtubules. Regulates the kinesin-mediated axonal transport of mitochondria to nerve terminals along microtubules during hypoxia. Participates in the translocation of TRAK2/GRIF1 from the cytoplasm to the mitochondrion. Also plays a role in steroidogenesis through maintenance of mitochondrial abundance and morphology (By similarity). Plays an inhibitory role during neocortex development by regulating mitochondrial morphology, distribution and motility in neocortical neurons (By similarity)

Product:

1mg/ml in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

Molecular Weight:

~25.4 kDa

Swiss-Prot:

Q8TDB4

Purification&Purity:

The antibody was purified by immunogen affinity chromatography.

Applications:

WB (1/500 - 1/1000)

Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Recognizes endogenous levels of MGARP protein

DATA:

Western blot analysis of Tetranectin expression in mouse heart (A), rat heart (B) whole cell lysates.

Immunohistochemical analysis of Tetranectin staining in human kidney formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Immunofluorescent analysis of Tetranectin staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151