



SPATA2 polyclonal antibody

Catalog: BS35649

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Spermatogenesis-associated protein 2: Bridging factor that mediates the recruitment of CYLD to the LUBAC complex, thereby regulating TNF-induced necroptosis . Acts as a direct binding intermediate that bridges RNF31/HOIP, the catalytic subunit of the LUBAC complex, and the deubiquitinase (CYLD), thereby recruiting CYLD to the TNF-R1 signaling complex (TNF-RSC) . Required to activate the 'Met-1'- (linear) and 'Lys-63'-linked deubiquitinase activities of CYLD . Controls the kinase activity of RIPK1 and TNF-induced necroptosis by promoting 'Met-1'-linked deubiquitination of RIPK1 by CYLD (By similarity)

Product:

1mg/ml in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.

Molecular Weight:

~58.4 kDa

Swiss-Prot:

Q9UM82

Purification&Purity:

The antibody was purified by immunogen affinity chromatography.

Applications:

WB (1/500 - 1/1000), IHC (1/50 - 1/200)

Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long

term. Avoid freeze-thaw cycles.

Specificity:

Recognizes endogenous levels of SPATA2 protein

DATA:

Western blot analysis of Aldose Reductase expression in HEK293T (A), Hela (B), mouse testis (C), rat testis (D) whole cell lysates.

Immunohistochemical analysis of Aldose Reductase staining in human colon cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Immunofluorescent analysis of Aldose Reductase staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151