

CDC40 Recombinant Rabbit mAb

Catalog: BS50159

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Pre-mRNA splicing occurs in two sequential transesterification steps. The protein encoded by this gene is found to be essential for the catalytic step II in pre-mRNA splicing process. It is found in the spliceosome, and contains seven WD repeats, which function in protein-protein interactions. This protein has a sequence similarity to yeast Prp17 protein, which functions in two different cellular processes: pre-mRNA splicing and cell cycle progression. It suggests that this protein may play a role in cell cycle progression. [provided by RefSeq, Jul 2008]

Product:

1mg/ml in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA.

Molecular Weight:

75 kDa

Swiss-Prot:

O60508

Purification&Purity:

Affinity Purification

Applications:

WB: 1:1000 IHC-P: 1:100 ICC/IF: 1:100-1:200 FC: 1:20-1:100


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.


Specificity:

The antibody detects endogenous levels of it.

DATA:



Western blot analysis of extracts from K562, HeLa cells and Mouse spleen tissue using it at 1:1000.



Immunofluorescence analysis of HeLa cells labelling CDC5L with it.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151